

**U. PORTO**



FACULDADE DE MEDICINA  
UNIVERSIDADE DO PORTO

**MESTRADO INTEGRADO EM MEDICINA**

---

2017/2018

João Pedro Rosa de Oliveira

# Vascular Access for Elderly Hemodialysis Patients

Acessos Vasculares em Doentes Idosos em Hemodiálise

março, 2018

FMUP

João Pedro Rosa de Oliveira

## **Vascular Access for Elderly Hemodialysis Patients**

**Acessos Vasculares em Doentes Idosos em Hemodiálise**

**Mestrado Integrado em Medicina**

**Área: Nefrologia**

**Tipologia: Artigo de Revisão**

**Trabalho efetuado sob a Orientação de:**

**Doutor Luís Alexandre de Castilho Silva Coentrão**

**Trabalho organizado de acordo com as normas da revista:**

**Portuguese Journal of Nephrology and Hypertension**

março, 2018

**FMUP**

Eu, João Pedro Rosa de Oliveira, abaixo assinado, nº mecanográfico 201102571, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 21/03/2018

Assinatura conforme cartão de identificação:

João Pedro Rosa de Oliveira



NOME

João Pedro Rosa de Oliveira

NÚMERO DE ESTUDANTE

E-MAIL

2011 02571 joãpedro2oliveira93@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Medicina - Nefrologia

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

vascular access for Elderly Hemodialysis Patients

ORIENTADOR

Dr. Luís Alexandre de Castilho Silva Coentrão

COORDINADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

|   |                                     |
|---|-------------------------------------|
| É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.   | <input checked="" type="checkbox"/> |
| É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE. | <input type="checkbox"/>            |
| DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.  | <input type="checkbox"/>            |

Faculdade de Medicina da Universidade do Porto, 20/3/18

Assinatura conforme cartão de identificação:

João Pedro Rosa de Oliveira

**Full Title:**

# **Vascular Access for Elderly Hemodialysis Patients**

**Authors' names and institutional addresses:**

João Pedro Oliveira, Luís Coentrão

Department of medicine, Faculty of Medicine, University of Porto, Porto Portugal

**Author:**

João Pedro Rosa de Oliveira

**Address:** Ruas Cidade de Amadora nº10 2ºD 2855-116 Corroios

**Email:** [joapedroroliveira93@gmail.com](mailto:joapedroroliveira93@gmail.com)

# **Abstract**

**Introduction:** Arteriovenous Fistulas have been massively considered as the gold-standard vascular access in Hemodialysis starting patients. Although data shows AVFs as the best method overall, in older patients some contradiction has been noted. Therefore, rises the question, are still AVFs the best method for elderly patient's treatment? And if so, why?

**Methods:** Creation of this paper was based on data gathering. Through pubmed's search engine, we looked for English written articles discussing the topic of Hemodialysis vascular access in the elderly. Articles on access comorbidities and outcomes were also included. Studies prior to the year 2000 were excluded.

**Results:** The association between old age and high cardio-vascular comorbidity rates is linked to poorer fistula maturation. Poor out-comes and low life-expectancy leads to the belief that fistulas may not be recommended for elder patients. However, contradictory studies make it impossible to come to a proper conclusion. Still, most authors seem to agree that proper patient guidance during and prior to Hemodialysis treatment transition, may lead to better access selection and better fistula outcome. Vascular examination with ultrasound technics and well timed vascular access creation (6-9 months before treatment) are also associated to better fistula success. Also, proximal fistula creation (braquial-cephalic), was deemed better for older populations when compared to distal fistula creation, due to more significant patency.

**Conclusion:** Due to antithetical results, arteriovenous fistulas may not be disregarded as the primary access for HD treatment in the elder population, and fistula creation should still be carried-out in these patients. Nevertheless, patient treatment individualization is key and non-maturation risk factors should be searched and considered, as well as poor life-expectancy and bad vasculature.

**Key words:** Arteriovenous fistula, Elderly, Hemodialysis, Treatment failure, Vascular access.

## **Introduction**

Since 2003, the Fistula First Initiative has been one of the major deciding factors while choosing the vascular access for initiating Hemodialysis (HD) treatment <sup>(1)</sup>. Since then most guidelines suggest Arteriovenous Fistulas (AVF) as the gold-standard access for all patients beginning HD worldwide <sup>(2)</sup>. Despite the increased numbers of Fistulas created, elder patients treated with HD continue to have high mortality rates, independently of vascular access<sup>(3)</sup>. Relation between age and mortality is unequivocal, still questions regarding current vascular access methods in elders have since been raised <sup>(4)</sup>.

## **Methods**

A diverse literature search was made using Pubmed's (U.S. National Library of Medicine), in order to identify relevant articles on the topic of vascular access in the elderly beginning hemodialysis. Arteriovenous fistula and arteriovenous graft were used as keywords, as well as hemodialysis, elderly and chronic kidney failure. Studies on access and technic comorbidities were also included. English written articles were preferred due to linguistic barriers. Studies prior to the year 2000 were excluded from the selection. Moreover, copies of the papers were acquired on electronic format from the retrieved journals, in order to enhance the search for information on the topic.

## **Vascular Access Choice**

When choosing between the available vascular access for patients starting HD, AVFs are the preferable option. Fistulas have been associated with a better life-span, lower rates of hospitalization and infection in comparison to arteriovenous grafts (AVG) and central venous catheters (CVC) <sup>(5, 6)</sup>. The difference in methods, their contradictions and indications have long been a known fact and because of that AVFs have been considered as the more favourable method and gold-standard treatment <sup>(2)</sup>. However, recent studies showed a lack of benefit in fistula creation in the elderly population, which contradicts HD treatment guidelines<sup>(7-10)</sup>. DeSilva et al, compared data from a cohort of 115,425 HD patients > 67 years old and failed to show a statistical difference between fistula and AVG usage in patients older than 80 <sup>(7)</sup>. Other studies even claim that “high

risk patients may benefit from permanent central vein catheters” or “long term catheters”<sup>(11, 12)</sup>. Disparity in results and contradictions to the Fistula First Program, creates doubt in clinical decision, specifically in this group of patients. So, we may question, what’s the preferred method for elders beginning HD?

The importance behind likewise questions regards not only the well-fare and outcome of patients, but also the proper and beneficial management of hospital resources, that are in high demand due to the increase in the HD population<sup>(1)</sup>. Moreover, divergent data questions the reliability of AVFs as the preferred treatment and the Fistula First Program itself. Furthermore, current guidelines lack orientation on the matter, suggesting only that individualized AV planning may lead to a better outcome in elder patients, without a specific answer on the preferable method for the older patients<sup>(13)</sup>.

One thing is certain, lack of success in the elderly is mostly associated to an increased AVF maturation failure<sup>(8, 13, 14)</sup>.

### **AVF failure**

Many variables have been correlated to an increase in Fistula failure. Metabolic syndrome, Diabetes<sup>(15)</sup>, Atherosclerosis, thrombosis, female sex<sup>(16)</sup> and black race<sup>(17)</sup>, make some of the most important<sup>(18-20)</sup>. It isn’t hard to relate most of the previously referred criteria to old age. Older people are largely associated with increased levels of cardiovascular comorbidities, making it easy to justify the difficulty in AVF creation in this group. Additionally, older age seems to increase AVF failure independently of other risk factors. Mauricio Monroy-Cuadros et al, studying a cohort of 831 patients after AVF creation, disclosed that the risk of AVF failure is similar, independently, for age, Diabetes, Hypertension and peripheral vascular disease (RR: 2.4-2.5). More than that, results show a 16 times risk increase in patients with low initial intra-access blood flow, that is also linked to age related comorbidities like atherosclerosis and Peripheral vascular disease(PVD)<sup>(21)</sup>.

PVD and low initial intra-access blood flow, are highly associated to peripheral hypoperfusion. As previously referred, poor peripheral vasculature has the highest risk of AVF non-maturation and higher chance of thrombosis<sup>(21)</sup>. Regarding this information, a proper vascular examination is required in HD starting patients, especially in the elderly<sup>(13)</sup>. Thus, recommendation for ultrasound vascular exams to be carried on in these



patients. Artery and Venous calibre should be considered, such as flow rate and any vascular turbulence due to atherosclerosis, since any of previous factors may question the maturation of the fistula.

Due to vascular abnormalities and deficient patency, in the elder vasculature, location of the vascular access is also key for a proper maturation. Most researchers seem to agree that proximal fistula is related to better success rates. This might be attributed to the higher calibre and blood flow of the vessels. So, according to the previous information braquiocephalic fistula are preferred to radiocephalic. Braquiocephalic access has a better life-span and a lower failure percentage. Indeed studies show a better outcome when patients receive a braquiocephalic access<sup>(22, 23)</sup>. This association is specially well founded for older patients due to PVD<sup>(21)</sup>.

Certainly, the previous factors make the requirement for proper clinical evaluation in elders starting HD highly understandable and expected.

### **Proper care in access creation**

Thorough examination and overview are beneficial not only for the overall patient's well-being but for fistula creation as well. All HD patients seem to benefit from a prior well guided follow-up, which is essential for proper access creation<sup>(8, 10, 14, 24)</sup>.

Deciding when and if HD should start is also dependent on a good follow-up. A well-founded decision to begin treatment prior to terminal kidney disease may prevent life-threatening events and increase life-expectance as well as quality of life. It is also important to explain and discuss with patients the preferred method of treatment, including vascular access, since access outcome is dependent of the patient's cooperation and understanding.

Timing of access creation is also a fundamental factor in lowering fistula and graft failure rates. Proper timing may be the difference between fistula maturation and non-maturation, more so in elders. Studies show that too early access creation (> 12 months) may be prejudicial<sup>(25, 26)</sup>. Tammy Hod et al, relates a slight increase in Fistula maturation when access creation is started 6 to 9 months prior to HD<sup>(25)</sup>. However, due to the difficulty in predicting CKD evolution, most patients still don't start access creation in this

time range and continue to be subjected to fistulas with not enough time for proper maturation.

Overall recommendations suggest that AVF should always be considered, independently of age. However, individuality should also be considered, especially in the elderly, high comorbidity, low life-span patients. Some seem to agree that patients with less than 24 months of life-expectancy don't benefit from AVF creation, due to low maturation rates<sup>(27)</sup>. Still, one may question this statement due to the variability and difficulty in predicting life-expectancy.

## **Conclusion**

In conclusion, the lack of agreement on the subject makes it impossible to distinguish the best vascular access for the elderly, with the data available so far. Knowing this, it is essential that further, more specific study on the topic is conducted. However, the huge majority of authors agree that therapy should be individualized, due to the high comorbidities present and that AVFs, with the available data, should still be tried out as treatment for elder patients<sup>(21, 28-33)</sup>.

## **References**

1. Saran R, Li Y, Robinson B, Abbott KC, Agodoa LY, Ayanian J, et al. US Renal Data System 2015 Annual Data Report: Epidemiology of Kidney Disease in the United States. American journal of kidney diseases : the official journal of the National Kidney Foundation. 2016;67(3 Suppl 1):Svii, S1-305.
2. Clinical practice guidelines for vascular access. American journal of kidney diseases : the official journal of the National Kidney Foundation. 2006;48 Suppl 1:S248-73.
3. Coric A, Resic H, Celik D, Masnic F, Ajanovic S, Prohic N, et al. Mortality in hemodialysis patients over 65 years of age. Materia socio-medica. 2015;27(2):91-4.
4. Hicks CW, Canner JK, Arhuidese I, Zarkowsky DS, Qazi U, Reifsnyder T, et al. Mortality benefits of different hemodialysis access types are age dependent. Journal of vascular surgery. 2015;61(2):449-56.
5. Zhang JC, Al-Jaishi AA, Na Y, de Sa E, Moist LM. Association between vascular access type and patient mortality among elderly patients on hemodialysis in Canada. Hemodialysis international International Symposium on Home Hemodialysis. 2014;18(3):616-24.

6. Ferreira H, Coentrao L. Vascular access for elderly hemodialysis patients: what should we aim for? *The journal of vascular access*. 2016;17 Suppl 1:S38-41.
7. DeSilva RN, Patibandla BK, Vin Y, Narra A, Chawla V, Brown RS, et al. Fistula first is not always the best strategy for the elderly. *Journal of the American Society of Nephrology : JASN*. 2013;24(8):1297-304.
8. Moist LM, Lok CE, Vachharajani TJ, Xi W, AlJaishi A, Polkinghorne KR, et al. Optimal hemodialysis vascular access in the elderly patient. *Seminars in dialysis*. 2012;25(6):640-8.
9. Chan MR, Sanchez RJ, Young HN, Yevzlin AS. Vascular access outcomes in the elderly hemodialysis population: A USRDS study. *Seminars in dialysis*. 2007;20(6):606-10.
10. Nadeau-Fredette AC, Goupil R, Montreuil B, Carignan A, Leblanc M. Arteriovenous fistula for the 80 years and older patients on hemodialysis: is it worth it? *Hemodialysis international International Symposium on Home Hemodialysis*. 2013;17(4):594-601.
11. Vachharajani TJ, Moist LM, Glickman MH, Vazquez MA, Polkinghorne KR, Lok CE, et al. Elderly patients with CKD--dilemmas in dialysis therapy and vascular access. *Nature reviews Nephrology*. 2014;10(2):116-22.
12. Tordoir JH, Bode AS, van Loon MM. Preferred strategy for hemodialysis access creation in elderly patients. *European journal of vascular and endovascular surgery : the official journal of the European Society for Vascular Surgery*. 2015;49(6):738-43.
13. Sequeira A, Naljayan M, Vachharajani TJ. Vascular Access Guidelines: Summary, Rationale, and Controversies. *Techniques in vascular and interventional radiology*. 2017;20(1):2-8.
14. Leimbach T, Kron J, Czerny J, Urbach B, Aign S, Kron S. Hemodialysis in patients over 80 years. *Nephron*. 2015;129(3):214-8.
15. Patibandla BK, Narra A, Desilva R, Chawla V, Vin Y, Brown RS, et al. Disparities in arteriovenous fistula placement in older hemodialysis patients. *Hemodialysis international International Symposium on Home Hemodialysis*. 2014;18(1):118-26.
16. Goldfarb-Rumyantzev AS, Rout P. Characteristics of elderly patients with diabetes and end-stage renal disease. *Seminars in dialysis*. 2010;23(2):185-90.
17. Drew DA, Lok CE, Cohen JT, Wagner M, Tangri N, Weiner DE. Vascular access choice in incident hemodialysis patients: a decision analysis. *Journal of the American Society of Nephrology : JASN*. 2015;26(1):183-91.
18. Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, et al. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet (London, England)*. 2010;376(9739):419-30.

19. Hayakawa K, Miyakawa S, Hoshinaga K, Hata K, Marumo K, Hata M. The effect of patient age and other factors on the maintenance of permanent hemodialysis vascular access. *Therapeutic apheresis and dialysis : official peer-reviewed journal of the International Society for Apheresis, the Japanese Society for Apheresis, the Japanese Society for Dialysis Therapy*. 2007;11(1):36-41.
20. Woo K, Gascue L, Goldman DP, Romley JA. Variations in outcomes of hemodialysis vascular access by race/ethnicity in the elderly. *Journal of vascular surgery*. 2017;65(3):783-92.e4.
21. Lilly MP, Lynch JR, Wish JB, Huff ED, Chen SC, Armistead NC, et al. Prevalence of arteriovenous fistulas in incident hemodialysis patients: correlation with patient factors that may be associated with maturation failure. *American journal of kidney diseases : the official journal of the National Kidney Foundation*. 2012;59(4):541-9.
22. Monroy-Cuadros M, Yilmaz S, Salazar-Banuelos A, Doig C. Risk factors associated with patency loss of hemodialysis vascular access within 6 months. *Clinical journal of the American Society of Nephrology : CJASN*. 2010;5(10):1787-92.
23. Misskey J, Faulds J, Sidhu R, Baxter K, Gagnon J, Hsiang Y. An age-based comparison of fistula location, patency, and maturation for elderly renal failure patients. *Journal of vascular surgery*. 2017.
24. Sladoje-Martinovic B, Mikolasevic I, Bubic I, Racki S, Orlic L. Survival of chronic hemodialysis patients over 80 years of age. *Clinical interventions in aging*. 2014;9:689-96.
25. Hod T, Patibandla BK, Vin Y, Brown RS, Goldfarb-Rumyantzev AS. Arteriovenous fistula placement in the elderly: when is the optimal time? *Journal of the American Society of Nephrology : JASN*. 2015;26(2):448-56.
26. Woo K, Goldman DP, Romley JA. Early Failure of Dialysis Access among the Elderly in the Era of Fistula First. *Clinical journal of the American Society of Nephrology : CJASN*. 2015;10(10):1791-8.
27. Watorek E, Golebiowski T, Kusztal M, Letachowicz K, Letachowicz W, Augustyniak Bartosik H, et al. Creation of arteriovenous fistulae for hemodialysis in octogenarians. *Hemodialysis international International Symposium on Home Hemodialysis*. 2014;18(1):113-7.
28. DeSilva RN, Sandhu GS, Garg J, Goldfarb-Rumyantzev AS. Association between initial type of hemodialysis access used in the elderly and mortality. *Hemodialysis international International Symposium on Home Hemodialysis*. 2012;16(2):233-41.
29. Renaud CJ, Pei JH, Lee EJ, Robless PA, Vathsala A. Comparative outcomes of primary autogenous fistulas in elderly, multiethnic Asian hemodialysis patients. *Journal of vascular surgery*. 2012;56(2):433-9.
30. Palmes D, Kebschull L, Schaefer RM, Pelster F, Konner K. Perforating vein fistula is superior to forearm fistula in elderly haemodialysis patients with diabetes and arterial hypertension. *Nephrology, dialysis, transplantation : official publication of the*

European Dialysis and Transplant Association - European Renal Association.  
2011;26(10):3309-14.

31. Morsy M, Betal D, Nelson S, Malete H, Whitmore A, Chemla E. Pre-emptive angioaccess for haemodialysis in the elderly. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2011;26(11):3666-70.

32. Swindlehurst N, Swindlehurst A, Lumbair H, Rebollo Mesa I, Mamode N, Cacciola R, et al. Vascular access for hemodialysis in the elderly. *Journal of vascular surgery*. 2011;53(4):1039-43.

33. Bessias N, Paraskevas KI, Tziviskou E, Andrikopoulos V. Vascular access in elderly patients with end-stage renal disease. *International urology and nephrology*. 2008;40(4):1133-42.



ISSN 0872-0169  
[versão impressa]

## INSTRUÇÕES AOS AUTORES

- [Escopo e política](#)
- [Forma e preparação de manuscritos](#)

### Escopo e política

A Revista Portuguesa de Nefrologia e Hipertensão é o órgão oficial da Sociedade Portuguesa de Nefrologia e tem publicação trimestral, para além de Suplementos sobre temas seleccionados ao critério do Editor.

A Revista publica artigos sobre tópicos clínicos ou laboratoriais relevantes para nefrologia, diálise, transplantação renal e hipertensão. Trabalhos que versem sobre imunologia básica, fisiologia, genética e epidemiologia serão aceites se relacionados com aspectos renais.

A Revista Portuguesa de Nefrologia e Hipertensão publica Editoriais e Revisões abrangentes (apenas por convite), Artigos Originais, Comunicações Breves, Notas Técnicas, Casos Clínicos e Cartas ao Editor. Os artigos são submetidos e publicados apenas na língua inglesa. Os artigos submetidos devem ser acompanhados de carta, assinada por todos os autores, declarando a cedência dos direitos de propriedade a favor da Revista Portuguesa de Nefrologia e Hipertensão e a originalidade do trabalho, que não pode ter já sido publicado nem estar em processo de avaliação para publicação noutra revista.

Os manuscritos devem ser preparados em conformidade com os Requisitos Uniformes para Manuscritos Submetidos a Revistas Biomédicas, produzidos pelo ICMJE (International Committee of Medical Journal Editors), que poderão ser consultados em <http://www.icmje.org>. Todos os manuscritos recebidos são submetidos ao processo de revisão por pares.

### PROCESSO DE REVISÃO E PUBLICAÇÃO

Cada artigo recebido é imediatamente submetido a avaliação editorial, após o que é enviado a três revisores, especialistas na área, que desconhecem a autoria do trabalho em causa. O Editor envia todos os esforços na coordenação deste processo de modo a que se chegue a uma decisão no prazo máximo de 8 semanas, conseguindo, assim, que os trabalhos sejam publicados no número da Revista que se segue à sua aceitação. Os artigos que não correspondam aos padrões científicos da Revista Portuguesa de Nefrologia e Hipertensão poderão ser rejeitados pelo Editor sem recurso a outras revisões.

### Forma e preparação de manuscritos

O manuscrito, digitado em formato A4 e em espaço duplo, deve ser submetido ao Editor por correio electrónico para:  
[fcarrera@mail.telepac.pt](mailto:fcarrera@mail.telepac.pt)

**Primeira página:** Deve conter o título do artigo e os nomes e apelidos de todos os autores, bem como o nome e endereço da instituição onde



o trabalho foi desenvolvido; no caso de várias instituições, estas devem ser identificadas com algarismos e a respectiva correspondência aos autores. Nome completo e morada do autor correspondente. Com início nesta, todas as páginas devem ser numeradas consecutivamente.

Na segunda página deverá constar novamente o título do artigo.

**Resumo:** Deverá conter até 300 palavras, sem abreviaturas nem acrónimos.

**Palavras-Chave:** Num máximo de 6, por ordem alfabética e, se possível, de acordo com a lista de Medical Subject Headings (Índex Medicus).

**Texto:** A ordem do texto deve ser a seguinte: Introdução, Material e Métodos (algum método estatístico utilizado deve ser detalhado nesta secção), Resultados, Discussão, Agradecimentos, Referências, Quadros, Legendas e Figuras.

**Quadros:** A referência aos quadros deve ser feita, por ordem de entrada no texto, entre parêntesis [e.g., (Quadro I)]. Cada quadro deve constar de uma folha separada, ter numeração romana e sequencial, bem como um título informativo do seu conteúdo.

**Figuras:** As figuras devem ser identificadas com numeração árabe e referidas, por ordem de entrada no texto, entre parêntesis [e.g., (Fig. 2)]. Uma figura que tenha sido publicada anteriormente deve ser acompanhada de cópia da autorização para re-publicação por parte da entidade detentora dos seus direitos de propriedade e reconhecida a respectiva fonte

**Referências Bibliográficas:** Devem ser numeradas e ordenadas segundo a ordem de citação no texto. As referências devem citar todos os autores (apelido e iniciais), quando até seis; acima deste número, devem citar-se os três primeiros seguido de et al. Aos nomes dos autores, segue-se o título do artigo, a abreviatura do nome da revista (segundo o estilo usado no Index Medicus), ano de publicação, número do volume, e números da primeira e última páginas. Referências a livros, devem incluir o nome do capítulo, nome do livro, local de publicação, editora, ano e páginas relevantes.

Exemplos:

1. Revista

Carrera F, Oliveira L, Maia P, Mendes T, Ferreira C. The efficacy of intravenous darbepoetin alfa administered once every 2 weeks in chronic kidney disease patients on haemodialysis. *Nephrol Dial Transplant* 2006;21:2846-2850. [SEP]

2. Livro de um único autor

Botella García J. *Manual de Nefrología Clínica*. 1st ed. Barcelona: Masson SA, 2003: 209.

3. Capítulo em livro

Carrera F, Frazão JM. Thromboembolism and atheroembolic disease of the kidney. In: Schena, FP, ed. *Nephrology*. London: McGraw Hill International, 2001: 355-359.

**Última página:** Nesta página devem constar os apoios e patrocínios (bolsas, equipamento, fármacos, etc.) concedidos. Os possíveis conflitos de interesses, no que se refere, por exemplo, a participação accionista, concessão de bolsas, pagamento de consultoria ou honorários pela indústria farmacêutica ou outra, devem ser explicitamente declarados em conformidade com as recomendações

ICMJE II:D1 sobre compromisso de autores e II:D2 sobre apoio a projectos.

[[Home](#)] [[Sobre esta revista](#)] [[Corpo Editorial](#)] [[Assinaturas](#)]

---

© 2012 *Publicações Ciência e Vida, Lda.*

*Praça Cidade de Odivelas, Lote 4, Loja Dta.  
2675 – 639 Odivelas*

*Tel. + 351 21 478 78 50*

*E-mail: [pub@cienciaevida.pt](mailto:pub@cienciaevida.pt)*